

Boerhaave Syndrome

Mitra Ahadi¹, Negin Masoudifar^{2*}

¹ Gastroenterologist, Department of Internal Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

² General Practitioner, Mashhad University of Medical Sciences, Mashhad, Iran

ARTICLE INFO

Article type:
Review Article

Article history:
Received: 24 Sep 2017
Revised: 18 Oct 2017
Accepted: 28 Oct 2017

Keywords:
Boerhaave Syndrome
Early Diagnosis
Esophageal Perforation

ABSTRACT

Boerhaave syndrome (BS) is an uncommon and life-threatening disorder and is defined as spontaneous transmural esophageal perforation, which typically occurs after a projectile vomiting. The prognosis is dependent on early detection of the disease and appropriate management within 12 h of perforation. The typical presentations of BS are vomiting, subcutaneous emphysema, and pain in the lower chest. However, severe symptoms rarely happen, and about one-third of all patients present with atypical clinical features. Therefore, BS should be suspected in any patient presenting with sudden thoracoabdominal pain with a history of vomiting. Chest X-ray is the most helpful diagnostic tool, and computed tomography scan is applied for further evaluations. Although surgical treatment is an important option for many patients, a less invasive procedure with or without endoscopic stent or internal or external drains, with respect to patient's clinical condition, is preferable.

► Please cite this paper as:

Ahadi M, Masoudifar N. Boerhaave Syndrome. J Cardiothorac Med. 2017; 5(4): 208-212.

Introduction

Boerhaave syndrome (BS) was first described by a Dutch physician and anatomist, Hermann Boerhaave, in 1724. His patient was a 50-year-old Dutch grand admiral, Baron Janvan Wassenaer, who died in 1723 after one hour of self-induced vomiting, which led to esophageal perforation. BS is one of the most life threatening diseases of the gastrointestinal tract with high mortality and morbidity rates (1-3).

The pathophysiology of BS involves a transmural esophageal perforation ensuing a sudden rise in intraluminal esophageal pressure due to vomiting against a closed glottis and failure of relaxation of the cricopharyngeal. The most common anatomical location of the rupture is at the left posterolateral wall of the distal esophagus, 2-3 cm proximal to the gastroesophageal junction. Moreover, the second common location of rupture is at the subdiaphragmatic or upper thoracic esophageal segments (4-8).

This condition usually occurs following excessive alcohol consumption or overeating, both of which can lead to vomiting. Because BS is an uncommon condition, misdiagnosis can

result in mediastinitis, sepsis and even shock. Timely diagnosis and effective management within 24 h reduce mortality rate by approximately 30%.

Without adequate treatment, BS survival is in days and the mortality rate approximates to 100% (9-13).

There are multiple factors deteriorating BS mortality rate, such as difficulty in accessing the esophagus, lack of a strong serosal layer, abnormality in blood supply, and proximity to the vital structures. Generally, BS mortality rate is approximately 30%, which is mainly due to subsequent infections including mediastinitis, pneumonitis, pericarditis, or empyema (14-16).

Epidemiology

According to the literature, this disorder is reported in all races. The annual incidence rate of esophageal perforations is low, approximately 3.1 per 1,000,000 individuals, and 15% of all cases have BS. Globally, BS is significantly prevalent in males, with male to female ratio of 2:1 to 5:1. BS is more common in the patients

*Corresponding author: Negin Masoudifar, Mashhad University of Medical Sciences, Mashhad, Iran. Tel: +989155254128; Email: masoudifar.negin@gmail.com

© 2017 mums.ac.ir All rights reserved.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

aged between 50 and 70 years. Although several studies demonstrated that 80% of all patients are middle-aged males, this syndrome is reported in neonates and in people older than 90 years, as well (17-21).

Clinical Manifestations

BS typically presents with a history of nausea and vomiting followed by a sudden onset of severe lower chest and epigastric pain. The pain may radiate to the back or left shoulder and often worsens after eating. Another common manifestation is shortness of breath due to pleurisy or a pleural effusion. Mackler's triad denotes the classic presentations of BS including vomiting, lower chest pain, and subcutaneous emphysema. Additionally, atypical findings of BS include hoarseness due to the involvement of recurrent laryngeal nerve and neck or upper chest pain in patients with cervical esophagus perforation (23-27).

Imaging and work-up

The diagnosis of BS relies on radiographic evidence.

Plain radiograph

Chest X-ray (CXR) findings are often non-specific and can be observed on a posteroanterior and lateral views. CXR findings of BS include pneumomediastinum, subcutaneous emphysema, left pleural effusion, and left pneumothorax. Left pleural infusion is one of the most common findings, and about 10% of CXRs are normal.

In 20% of patients Naclerio's V sign, which is a focal and sharply margined region of paraspinal radiolucency on the left side immediately above the diaphragm (Figure 1).




Figure 1. The Naclerio's V sign seen on the frontal radiographs of the chest as a V-shaped air lucency in the left lower mediastinal segment

Esophagography

Esophagography is an imaging technique that uses water-soluble contrast agents and often demonstrates a contrast leakage at supradiaphragmatic level in most esophageal perforation cases (Figure 2). Water-soluble contrast is utilized to prevent barium-related severe mediastinitis in patients with esophageal perforation, hence reduced morbidity and mortality. More than 20% of patients have reported false negative results; therefore, if clinical suspicion remains, Esophagography should be repeated after 4-6 h (27-30).

CT scan

An unenhanced CT scan reveals the presence of typical localization and periesophageal air collections indicating esophageal perforation. Post-contrast CT scan may show direct contrast extravasation and thickened esophageal walls. Other reported findings include pneumothorax,




Figure 2. Plain chest radiography with a water-soluble contrast swallow showing contrast leakage in a patient with Boerhaave syndrome




Figure 3. Mediastinal emphysema and extraluminal contrast in the left pleural cavity in patients with Boerhaave syndrome

pneumomediastinum, pleural effusion (usually left-sided), mediastinal fluid collections, and emphysema in the chest wall and neck (Figure 3). In comparison to CXR, CT scan better demonstrates the anatomical location and severity of perforation (31-34).

Endoscopy

There is controversy regarding the role of endoscopy in the diagnosis of BS. This procedure may raise the risk of extending the original perforation and forcing additional air through the perforation into the mediastinum. Endoscopy may be useful when a perforation is suspected, but radiographic findings are negative or it is impossible to use swallowing contrast agent (Figure 4) (35).

Treatment

Patients with suspected BS should be identified




Figure 4. Endoscopic view of a distal spontaneous perforation 24 h after the presentation of clinical symptoms. However, endoscopic appearance may suggest a period exceeding at least 36 to 48 h from the presentation of symptoms

at an early stage and immediately treated. Appropriate management consists of both conservative and surgical interventions. Choosing between conservative and aggressive treatment modalities is dependent on the extent of perforation, patient's medical condition, and delay in evaluation and presentation.

Medical care

In most patients, surgical management is the gold standard; nevertheless, non-invasive treatment might be convenient for the patients meeting the following criteria: A) The esophageal perforation is located in the mediastinum; B) There is a well drainage from the cavity into the esophagus; C) Patient presents with minimal symptoms; and D) There is minimal evidence of sepsis.

Conservative therapy includes intravenous fluids and broad-spectrum antibiotics (imipenem or cilastatin), nasogastric suction, proton pump inhibitors (PPIs), early use of nutritional support, and keeping the patient nil per os (NPO).

Surgical Care

If the diagnosis is made within the first 24 hours after perforation, primary repair is possible. Direct repair of the tear and drainage of the mediastinum and pleural cavity minimize the risk of further contamination and infection.

Surgical approach is determined by the location of esophageal perforation and surgeon's preference, and left thoracoabdominal incision is preferred. Omental flap is used for covering the primary closure. In addition, all the patients undergo laparotomy for gastrostomy and jejunostomy to help the drainage and nutrition.

Based on the study by de Schipper et al. (38), endoscopic procedure can be considered in certain patients. Endoscopy should be employed

for those patients diagnosed within 48 hours of the esophageal rupture without any signs of sepsis. However, if the patient presents signs of sepsis, thoracotomy with hemifundoplication and pleural/mediastinal drainage should be performed.

If diagnosis is made after 48 h of perforation, conservative therapy is recommended, and if septic profile is present, surgical treatment should be considered. According to the results of de Schipper et al., the survival rates of conservative, surgical, and endoscopic methods were 75%, 81%, and 100%, respectively (36-42).

Conclusion

The best outcome in BS is associated with early diagnosis and appropriate treatment. Due to the diverse range of presentations, the diagnosis is difficult and effective treatments are often delayed. Therefore, high index of suspicion for BS must be considered in any clinical condition in which the patient presents with a suspicious history with or without characteristic complaints.

Conflict of Interest

The authors declare no conflict of interest.

References

- Curci JJ, Horman MJ. Boerhaave's syndrome: the importance of early diagnosis and treatment. *Ann Surg.* 1976; 183:401-8.
- Walker WS, Cameron EW, Walbaum PR. Diagnosis and management of spontaneous transmural rupture of the oesophagus (Boerhaave's syndrome). *Br J Surg.* 1985; 72:204-7.
- Abbott OA, Mansour KA, Logan WD Jr, Hatcher CR Jr, Symbas PN. Atraumatic so-called "spontaneous" rupture of the esophagus. A review of 47 personal cases with comments on a new method of surgical therapy. *J Thorac Cardiovasc Surg.* 1970; 59:67-83.
- Teh E, Edwards J, Duffy J, Beggs D. Boerhaave's syndrome: a review of management and outcome. *Interact Cardiovasc Thorac Surg.* 2007; 6:640-3.
- Khan AZ, Strauss D, Mason RC. Boerhaave's syndrome: diagnosis and surgical management. *Surgeon.* 2007; 5:39-44.
- Soreide JA, Viste A. Esophageal perforation: diagnostic work-up and clinical decision-making in the first 24 hours. *Scand J Trauma Resusc Emerg Med.* 2011; 19:66-72.
- Spapen J, De Regt J, Nieboer K, Verfaillie G, Honoré PM, Spapen H. Boerhaave's syndrome: still a diagnostic and therapeutic challenge in the 21st century. *Case Rep Crit Care.* 2013; 2013:161286 .
- Wolfson D, Barkin JS. Treatment of boerhaave's syndrome. *Curr Treat Options Gastroenterol.* 2007; 10:71-7.
- Whyte RI. Boerhaave's syndrome. *N Engl J Med.* 2001; 344:139.
- Türüt H, Gulhan E, Adams PY, Cetin G. Successful conservative management of Boerhaave's syndrome with late presentation. *J Natl Med Assoc.* 2006; 98:1857-9.
- Witz M, Jedeikin R, Zager M, Shpitz B, Elyashiv A, Dinbar A. Spontaneous rupture of the distal oesophagus (Boerhaave's syndrome) with unusual clinical presentation of pneumoperitoneum. *Postgrad Med J.* 1984; 60:60-1.
- Kollmar O, Lindemann W, Richter S, Steffen I, Pistorius G, Schilling MK. Boerhaave's syndrome: primary repair vs. Esophageal resection--case reports and meta-analysis of the literature. *J Gastrointest Surg.* 2003; 7:726-34.
- Salo JA, Isolauri JO, Heikkilä LJ, Markkula HT, Heikkinen LO, Kivilaakso EO, et al. Management of delayed esophageal perforation with mediastinal sepsis. Esophagectomy or primary repair? *J Thorac Cardiovasc Surg.* 1993; 106:1088-91.
- Ryom P, Ravn JB, Penninga L, Schmidt S, Iversen MG, Skov-Olsen P, et al. Aetiology, treatment and mortality after oesophageal perforation in Denmark. *Dan Med Bull.* 2011; 58:A4267.
- Bhatia P, Fortin D, Inculet RI, Malthaner RA. Current concepts in the management of esophageal perforations: a twenty-seven year Canadian experience. *Ann Thorac Surg.* 2011; 92:209-15.
- Vidarsdottir H, Blondal S, Alfredsson H, Geirsson A, Gudbjartsson T. Oesophageal perforations in Iceland: a whole population study on incidence, aetiology and surgical outcome. *Thorac Cardiovasc Surg.* 2010; 58:476-80.
- Brinster CJ, Singhal S, Lee L, Marshall MB, Kaiser LR, Kucharzuk JC. Evolving options in the management of esophageal perforation. *Ann Thorac Surg.* 2004; 77:1475-83.
- Kish GF, Katske FA. A case of recurrent Boerhaave's syndrome. *W V Med J.* 1980. 76:27-30.
- Bladergroen MR, Lowe JE, Postlethwait RW. Diagnosis and recommended management of esophageal perforation and rupture. *Ann Thorac Surg.* 1986; 42:235-9.
- Restrepo CS, Lemos DF, Ocazionez D, Moncada R, Gimenez CR. Intramural hematoma of the esophagus: a pictorial essay. *Emerg Radiol.* 2008; 15:13-22.
- Ghanem N, Althoefer C, Springer O, Furtwängler A, Kotter E, Schäfer O, et al. Radiological findings in Boerhaave's syndrome. *Emerg Radiol.* 2003; 10:8-13.
- Shaker H, Elsayed H, Whittle I, Hussein S, Shackcloth M. The influence of the 'golden 24-h rule' on the prognosis of oesophageal perforation in the modern era. *Eur J Cardiothorac Surg.* 2010; 38:216-22.
- Derbes VJ, Mitchell RE Jr. Hermann Boerhaave's Atrocis, nec descripti prius, morbi historia, the first translation of the classic case report of rupture of the esophagus, with annotations. *Bull Med Libr Assoc.* 1955; 43:217-40.
- Buecker A, Wein BB, Neuerburg JM, Guenther RW. Esophageal perforation: comparison of use of aqueous and barium-containing contrast media. *Radiology.* 1997; 202:683-6.
- Rochford M, Kiernan TJ. Images in emergency medicine. Boerhaave's syndrome (spontaneous esophageal rupture). *Ann Emerg Med.* 2007; 49:746-77.
- Griffin SM, Lamb PJ, Shenfine J, Richardson DL, Karat D, Hayes N. Spontaneous rupture of the

- oesophagus. *Br J Surg*. 2008; 95:1115-20.
27. Mackler SA. Spontaneous rupture of the esophagus; an experimental and clinical study. *Surg Gynecol Obstet*. 1952; 95:345-56.
 28. Gimenez A, Franquet T, Erasmus JJ, Martinez S, Estrada P. Thoracic complications of esophageal disorders. *Radiographics*. 2002; 22:S247-58.
 29. Sinha R. Naclerio's V sign. *Radiology*. 2007; 245:296-7.
 30. Onyeka WO, Booth SJ. Boerhaave's syndrome presenting as tension pneumothorax. *J Accid Emerg Med*. 1999; 16:235-6.
 31. Sajith A, O'Donohue B, Roth RM, Khan RA. CT scan findings in oesophagogastric perforation after out of hospital cardiopulmonary resuscitation. *Emerg Med J*. 2008; 25:115-6.
 32. Exarhos DN, Malagari K, Tsatalou EG, Benakis SV, Peppas C, Kotanidou A, et al. Acute mediastinitis: spectrum of computed tomography findings. *Eur Radiol*. 2005; 15:1569-74.
 33. Huber-Lang M, Henne-Bruns D, Schmitz B, Wuerl P. Esophageal perforation: principles of diagnosis and surgical management. *Surg Today*. 2006; 36:332-40 .
 34. Young CA, Menias CO, Bhalla S, Prasad SR. CT features of esophageal emergencies. *Radiographics*. 2008; 28:1541-5.
 35. Arantes V, Campolina C, Valerio SH, de Sa RN, Toledo C, Ferrari TA, et al. Flexible esophagoscopy as a diagnostic tool for traumatic esophageal injuries. *J Trauma*. 2009; 66:1677-82.
 36. Bresadola V, Terrosu G, Favero A, Cattin F, Cherchi V, Adani GL, et al. Treatment of perforation in the healthy esophagus: analysis of 12 cases. *Langenbecks Arch Surg*. 2008; 393:135-40.
 37. Neel D, Davis EG, Farmer R, Richardson JD. Aggressive operative treatment for emetogenic rupture yields superior results. *Am Surg*. 2010; 76:865-8.
 38. de Schipper JP, Pull ter Gunne AF, Oostvogel HJ, van Laarhoven CJ. Spontaneous rupture of the oesophagus: Boerhaave's syndrome in 2008. Literature review and treatment algorithm. *Dig Surg*. 2009; 26:1-6.
 39. Vial CM, Whyte RI. Boerhaave's syndrome: diagnosis and treatment. *Surg Clin North Am*. 2005; 85:515-24.
 40. Ghanem N, Altehoefer C, Springer O, Furtwangler A, Kotter E, Schafer O, et al. Radiological findings in Boerhaave's syndrome. *Emerg Radiol*. 2003; 10:8-13.
 41. Abbas G, Schuchert MJ, Pettiford BL, Pennathur A, Landreneau J, Landreneau J, et al. Contemporaneous management of esophageal perforation. *Surgery*. 2009; 146:749-55.
 42. Kiernan PD, Rhee J, Collazo L, Hetrick V, Vaughan B, Graling P. Complete esophageal diversion: a simplified, easily reversible technique. *J Am Coll Surg*. 2005; 200:812.